



Available online at
ScienceDirect
www.sciencedirect.com

Elsevier Masson France
EM|consulte
www.em-consulte.com/en



Original article

Opening-wedge high tibial osteotomy with a secure bone allograft (Osteopure™) and locked plate fixation: Retrospective clinical and radiological evaluation of 69 knees after 7.5 years follow-up



G. Villatte^a, R. Erivan^a, P.-L. Fournier^a, B. Pereira^c, M. Galvin^a, S. Descamps^{a,b},
 S. Boisgard^{a,*}

^a Service d'orthopédie-traumatologie, CHU Montpied, 58, rue Montalembert, 63000 Clermont-Ferrand, France

^b Laboratoire C-Biosenss, EA 4676, Université d'Auvergne 1, 63000 Clermont-Ferrand, France

^c Service Biostatistique, CHU de Clermont-Ferrand, 58, rue Montalembert, 63000 Clermont-Ferrand, France

ARTICLE INFO

Article history:

Received 22 July 2015

Accepted 28 September 2015

Keywords:

Medial opening-wedge high tibial
 osteotomy
 Bone allograft
 Outcomes

ABSTRACT

Background: Medial opening-wedge high tibial osteotomy (OWHTO) provides reliable and long-lasting benefits, despite the wide range of wedge-filling and internal-fixation techniques used. The purpose of this work was to assess the clinical and radiological outcomes in a case-series of OWHTO performed using a secure bone allograft and locked plate fixation.

Hypothesis: The clinical and radiological outcomes of OWHTO with a high-safety bone allograft and locked plate fixation are similar to those reported in previous case-series studies.

Materials and methods: A single-centre retrospective design was used to study 69 knees in 64 patients with a mean age of 51.8 years (31–53 years) and a preoperative hip-knee-ankle (HKA) angle of 173° (165°–180°). The wedge was filled with secure Osteopure™ bone allograft and fixation was achieved using an Integra Surfix® locked plate. Mean follow-up was 7.5 years (5–9.3 years). Clinical and functional outcomes were assessed by determining the IKS and KOOS-PS scores and recording complications related or unrelated to the allograft. The main criterion for assessing OWHTO survival was the time to revision surgery for symptom recurrence. Radiological assessment criteria were the HKA angle, tibial slope, patellar height, and osteoarthritis grade. GESTO criteria were used to evaluate the behaviour of the allograft.

Results: Of the 69 knees, 64 (92.8%) were re-evaluated. The survival rate after 9.3 years was 95% ± 2.7% (3 failures managed with arthroplasty). The functional IKS score improved significantly, by 20 points ($P < 0.001$). Mean increases of 7.8° for the HKA angle and 3.5° for tibial slope were recorded. Bone healing without radiological abnormalities was consistently achieved within 6 months. There were no complications related to the allograft (infections, allergies; local or systemic toxicity).

Discussion: The clinical, radiological, and safety outcomes documented in our study were similar to those reported in earlier work.

Level of evidence: IV, retrospective case-series study.

© 2015 Elsevier Masson SAS. All rights reserved.

1. Introduction

The use of surgery to treat knee osteoarthritis is expanding [1], and the target population is composed of increasingly young patients with long life expectancies. Knee osteoarthritis affects the medial tibiofemoral compartment in 70% of cases, usually in knees with varus malalignment.

Medial opening-wedge high tibial osteotomy (OWHTO) is a rational and validated procedure that provides lasting benefits, with good results in 85% of cases after 10 years [2,3]. The tibial wedge is usually filled. Autografts can be used but are associated with donor-site morbidity [4–8]. Consequently, researchers have developed a number of other options such as allografts, acrylic cement, and bone substitutes [9]. Filling materials must be biologically safe and capable of osteo-integration and healing. A specific locked plate designed to ensure rigid internal fixation used without filling the wedge has produced satisfactory outcomes [10,11].

The objective of this study was to assess outcomes in consecutive patients at a mean of 7.5 years after medial OWHTO with

* Corresponding author.

E-mail addresses: sboisgard@chu-clermontferrand.fr,
stephane.boisgard@wanadoo.fr (S. Boisgard).

implantation of the secure bone allograft Osteopure™ and locked plate fixation. The working hypotheses were that clinical and radiological outcomes with this technique were at least as good as those reported with other methods and that Osteopure™ was not associated with any specific complications.

2. Materials and methods

2.1. Study design

A retrospective single-centre design was used to study consecutive patients in whom medial OWHTO was carried out in our department between January 2004 and December 2007. After a mean of 7.5 years (5.2–9.4 years), patients underwent a clinical and radiological assessment performed by an independent observer.

The study protocol was approved by the ethics committee of the Clinical Investigations Centre (CECIC) of the Rhône-Alpes/Auvergne inter-region (IRB 00005921). Patients were informed about the study.

2.2. Study population

The study included 69 knees in 64 patients, 43 males and 21 females with a mean age of 51.8 years (range, 31–74 years). Mean preoperative body mass index (BMI) was $27.2 \pm 6.4 \text{ kg/m}^2$. The diagnosis was degenerative joint disease in all 69 knees. Radiological osteoarthritis severity according to the Ahlback system [12] was as follows: grade 1, 32%; grade 2, 53.5%; grade 3, 14.5%; and grade 4, 0%. Mean varus deformity was 173° (range, 165° – 180°). The mean Caton-Deschamps index of patellar height [13] was 0.90 (range, 0.5–1.25) and the mean tibial slope was 5.3° (range, 0° – 16°).

2.3. Operative technique

Osteopure™ was distributed by Ostéobanque d'Auvergne (Clermont Ferrand, France) as wedges (height, 10 or 15 mm; and slope, 6° or 10°) specifically designed for filling the defects created during OWHTO. The allogeneic bone was harvested during primary total hip arthroplasty procedures. Virus inactivation and sterilisation was achieved using a urea-based procedure and beta radiation carried out by OST Développement (Clermont Ferrand, France). This procedure was previously validated by the microbiology safety task force of the French Drug and Healthcare Safety Agency (AFSSAPS, accreditation # 99-08802B02), and bone allografts treated with this procedure were first used in humans in 1999.

Internal fixation was with the Surfix® locked plate (Integra, Saint Priest, France). Surfix® is a thick L-shaped plate specifically designed to ensure stability in all three planes after OWHTO. Threaded holes lock the screws into the plate.

The amount of correction needed was determined based on a weight-bearing evaluation of lower limb alignment. The target HKA angle was $183^\circ (\pm 3^\circ)$.

No immobilisation was used after surgery. A non-weightbearing rehabilitation programme was followed for 45 days. Prophylactic anticoagulant therapy was given for 2 months.

2.4. Assessment methods

Patients were required to have two clinical and radiological evaluations, one performed preoperatively and the other done at last follow-up by an independent observer. Revision surgery consisting in total knee arthroplasty (TKA) was the endpoint used to assess OWHTO survival. Functional outcomes were assessed by determining the Charnley functional class and IKS score preoperatively and at last follow-up, as well as the KOOS-PS (knee-related quality

Table 1

GESTO criteria for evaluating bone graft appearance [15].

Evaluation of the host bone: no change, sclerosis, lysis
Evaluation of the host bone/graft interface: surrounding line absent or present (size <25%, 25–50%, 50–75%, >75%, 100%; and thickness in millimetres)
Evaluation of the graft: sclerosis, fragmentation, migration, resorption (none, <25%, 25–50%, 50–75%, >75%, 100%)

of life) at last follow-up. All complications were recorded, including general complications (e.g., infection, intraoperative fracture, thrombo-embolism) and adverse events directly related to the bone allograft (contamination with infectious agents, allergy, acute or chronic toxicity).

Standard radiographs (weight-bearing anteroposterior and lateral views and skyline view with the knee flexed at 30°) and an assessment of lower-limb alignment were used to determine the following: tibiofemoral osteoarthritis grade in the Ahlback classification system, mechanical axis of the knee (HKA angle), tibial slope according to Dejour [14], Caton-Deschamps index of patellar height, and degree of bone healing. The appearance of the graft was assessed based on criteria developed by the Greffes Et Substituts Tissulaires en Orthopédie (GESTO) [15] (Table 1).

2.5. Statistical tests

Paired data (preoperative vs. postoperative values) were compared using Student's or Wilcoxon's test. Categorical variables in independent groups were compared by applying the chi-squared test or Fisher's exact test. Associations (multivariate analysis) between quantitative variables were assessed by computing Pearson's or Spearman's correlation coefficients. The Kaplan-Meier method was applied to assess data that were considered censored, namely, survival and revision surgery. Values of $P < 0.05$ were considered significant. All analyses were performed for a two-sided Type I risk of 5% using STATA V12 software (StataCorp, College Station, TX, USA).

3. Results

3.1. General results

After a mean follow-up of 7.5 years, 64 knees in 59 patients were assessed based on an in-person visit (91%) or on information from a telephone call and imaging studies (9%). Data were not available for 3 (4.7%) patients who were lost to follow-up and 2 (3.1%) who died.

OWHTO failed in 3 knees, after a mean of 5.25 ± 0.86 years. In all 3 cases, TKA was performed because of symptom recurrence and radiological osteoarthritis progression (HKA angle values at revision surgery: 179° , 189° , and 184° , respectively).

The Kaplan-Meier analysis showed a survival rate of $95\% \pm 2.7\%$ after 7.5 years (Fig. 1).

3.2. Complications

Intraoperative fracture of the lateral tibial plateau occurred in 3 (5.1%) cases and consistently healed within 3 months. This complication had no impact on the HKA angle at last follow-up (180° , 183° , and 184° , respectively). Deep vein thrombosis developed in 2 (3.4%) patients, both of whom achieved a full recovery with curative-dose anticoagulant therapy. No blood vessel or nerve injuries were recorded, and none of the patients experienced non-union.

There were no complications specifically related to the bone allograft (infection, allergy, local or systemic toxicity).




Fig. 1. Survival curve (revision with TKA as the endpoint).




Fig. 2. Changes over time in the distribution of patients according to Charnley class.

3.3. Clinical outcomes

At last follow-up, significant worsening ($P < 0.001$) of the Charnley class was noted, with a shift towards class C (Fig. 2). The total IKS score improved significantly ($P < 0.001$), from 132 (95% confidence interval [95%CI], 127–136) preoperatively to 151 (95%CI, 142–159) at last follow-up. The same significant differences were noted for the clinical and functional components of the score.

The mean KOOS-PS score at last follow-up was 38.8/100. The score result was considered good in 45% of cases, fair in 49%, and poor in 6%.

3.4. Radiological outcomes

The HKA angle target of $183^\circ \pm 3^\circ$ was achieved in nearly 70% of cases (mean HKA angle, 180.9° [176° – 189°]). Tibial slope increased significantly, by a mean of 3.5° ($P < 0.001$). The patellar index decreased significantly, by 0.19 ($P = 0.001$) (Table 2).

The osteoarthritis severity grade worsened in both the tibio-femoral and patellofemoral compartments (Fig. 3).

No evidence of allograft fragmentation was seen 6 months after surgery, when the density of the implant was either equal (82%) or higher (18%) than that of the surrounding bone. Evaluation of the GESTO criteria did not show any lines at the host bone-allograft

Table 2
Changes in mean values of radiological variables during the study.

	Preoperative values	After 7.5 years	p value
HKA angle ($^\circ$)	173.1	180.9	< 0.001
Tibial slope ($^\circ$)	5.3	8.8	< 0.001
Patellar index	0.90	0.71	= 0.001

HKA: mechanical axis of the lower limb.




Fig. 3. Changes over time in tibio-femoral osteoarthritis severity (Ahlbäck grade).

interface or changes in the host bone in contact with the implant. Finally, healing was achieved in all cases (Fig. 4).

3.5. Multivariate analysis

None of the clinical or radiological variables (including age, BMI, and HKA angle at last follow-up) correlated significantly with failure of the OWHTO procedure. The only significant association found was between worsening of tibiofemoral osteoarthritis severity and the clinical/functional IKS score at last follow-up ($P < 0.05$).

4. Discussion

This retrospective study of 64 knees managed consecutively with OWHTO then re-evaluated after a mean of 7.5 years shows that filling the wedge with bone allograft and using locked plate fixation produces similar outcomes to those reported previously [2,3,11,16–28] (Table 3).

The mean follow-up of 7.5 years in our population may seem too short, given that OWHTO outcomes are known to deteriorate over time, particularly after 10 to 15 years [2]. Furthermore, the loss of HKA angle correction over time (difference between the early post-operative value and the value at last follow-up) was not assessed, a fact that limits the validity of the analysis of the correction achieved.

4.1. Clinical and radiological outcomes

Medial OWHTO is a safe technique associated with fewer complications than lateral closing-wedge HTO. In particular, the frequency of nerve injury is low (0–8.3%) [2,29–32]. The data from our case-series confirm the low adverse event rate by documenting the absence of complications having noticeable effects on the patients.

The radiological outcomes were satisfactory at last follow-up. Overall, the HKA angle was returned to the normal range (mean, 181°) and was within the predefined target of $183^\circ \pm 3^\circ$ in 70% of patients. This slight undercorrection is probably ascribable to loss of correction over time (about 3° over 10 years on average) [2,33,34], although we did not evaluate this point. Another hypothesis is insufficient initial correction during surgery. Given this possibility, and in agreement with the conclusions of the 2008 SOFCOT symposium [2], we now target an HKA angle of 183° to 186° in the coronal plane.

In contrast to most of the previous studies [2,4,33,35], this tendency towards undercorrection (mean HKA angle, 181°) was not correlated with OWHTO survival (95% after 7.5 years), which was similar to that reported in recent work [2,3,11,16–28]. This apparent contradiction may be ascribable in part to the short follow-up. Similarly, the HKA angle did not correlate with the IKS score, which improved by 20 points.

Tibial slope increased by 3.5° and the patellar index decreased by 0.21 in our study, in keeping with previously reported radiological changes induced by OWHTO [2,36,37,38,39,40,41]. These changes




Fig. 4. Pre-operative long-leg radiograph showing 5° of varus. Intraoperative photograph of the Osteopure™ graft and locked plate fixation. Radiograph at last follow-up after 7.5 years.

Table 3
Survival rates of opening-wedge high tibial osteotomy in recent case-series studies.

Study	Year	Number of patients	Techniques	Follow-up (years)	Survival rate (%)
Dubrana et Sofcot [2]	2008	164	Opening/Subtraction	11	85
Hernigou et al. [25]	2010	53	Opening + β Tricalcium Phosphate wedges	10	88.7
DeMeo et al. [18]	2010	20	Opening	8	70
Saragaglia et al. [23]	2011	124	Opening + β Tricalcium Phosphate wedges	5	88.8
Shallberger et al. [26]	2011	71	Opening/Subtraction	10	74
				10	92
				15	71
Niinimäki et al. [27] (meta-analysis)	2012	3195	Opening/Subtraction	5	89
Bode et al. [28] Harris et al. [20] (meta-analysis)	2013	4557	Opening without filling Opening/Subtraction	10	73
				5	96
				5	92.4
				10	84.5
				20	72.3
Bonasia et al. [17]	2014	99	Opening	7.5	75.9
Duivenvoorden et al. [19]	2014	92	Opening/Subtraction	6	85
Our study	2015	75	Opening	7.5	95

occurred despite the special attention given to the posterior positioning of the plate and allograft.

4.2. Wedge filling using Osteopure™ bone allograft

Different schools of practice exist regarding the need for filling the opening wedge. Nevertheless, filling seems to improve the quality and speed of healing [11], irrespective of the type of fixation plate [42].

Several materials can be used to fill the wedge. Autografting is the reference standard [42–44] based on good healing potential and low risk of infection. However, non-negligible donor site morbidity occurs (10–25% of complications) [4–8]. Many synthetic bone substitutes are available (e.g., calcium phosphate ceramics and calcium phosphate cements) [9]. They produce good healing rates [23,42,45,46], although residual traces of substitute often persist in the long term.

Secure bone allografts combine the theoretical advantages of the various materials (no donor-site morbidity, biological safety, good-quality healing, and restoration of the normal bone stock). The evaluation of GESTO criteria in this case-series confirmed these

advantages: the healing and resorption rates were similar to those seen with autografts, with no allograft-related morbidity or specific complications (in particular, no infections). These findings are in accordance with earlier reports [42–44,47–50].

Finally, the use of this high-safety bone allograft ensures the availability of a good bone stock in the event of subsequent revision surgery for TKA.

5. Conclusion

Medial OWHTO with a high-safety bone allograft and locking plate fixation is a safe and reproducible technique that produces similar outcomes to those of other procedures for managing knee osteoarthritis with varus malalignment in younger patients.

Disclosure of interest

S.B. and S.D. are members of the scientific committee of Ostéobanque d'Auvergne.

The other authors declare that they have no competing interest.

References

- [1] Singh JA. Epidemiology of knee and hip arthroplasty: a systematic review. *Open Orthop J* 2011;5:80–5.
- [2] Dubrana F, Lecerf G, Nguyen-Khanh J-P, Menard R, Ardouin L, Gibon Y, et al. Tibial valgus osteotomy. *Rev Chir Orthop Rep Appar Mot* 2008;94:S2–21.
- [3] Brouwer RW, Huizinga MR, Duivenvoorden T, van Raaij TM, Verhagen AP, Bierma-Zeinstra SMA, et al. Osteotomy for treating knee osteoarthritis. *Cochrane Database Syst Rev* 2014;12:CD004019.
- [4] Ahlmann E, Patzakis M, Roidis N, Shepherd L, Holtom P. Comparison of anterior and posterior iliac crest bone grafts in terms of harvest-site morbidity and functional outcomes. *J Bone Joint Surg Am* 2002;84:716–20.
- [5] Fowler BL, Dall BE, Rowe DE. Complications associated with harvesting autogenous iliac bone graft. *Am J Orthop* 1995;24:895–903.
- [6] Palmer W, Crawford-Sykes A, Rose REC. Donor site morbidity following iliac crest bone graft. *West Indian Med J* 2008;57:490–2.
- [7] Schaaf H, Lendeckel S, Howaldt H-P, Streckbein P. Donor site morbidity after bone harvesting from the anterior iliac crest. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2010;109:52–8.
- [8] Dimitriou R, Mataliotakis GI, Angoules AG, Kanakaris NK, Giannoudis PV. Complications following autologous bone graft harvesting from the iliac crest and using the RIA: a systematic review. *Injury* 2011;42(Suppl 2):S3–15.
- [9] Mainard D, Gouin F, Chauveaux D, Rosset P, Schwartz C. Les substituts de l'os, du cartilage et du ménisque en 2011 - SOFROT. Paris: Romillat; 2011.
- [10] Brinkman J-M, Lobenhoffer P, Agneskirchner JD, Staubli AE, Wymenga AB, van Heerwaarden RJ. Osteotomies around the knee: patient selection, stability of fixation and bone healing in high tibial osteotomies. *J Bone Joint Surg Br* 2008;90:1548–57.
- [11] Brosset T, Pasquier G, Migaud H, Gougeon F. Opening wedge high tibial osteotomy performed without filling the defect but with locking plate fixation (TomoFix™) and early weight-bearing: prospective evaluation of bone union, precision and maintenance of correction in 51 cases. *Orthop Traumatol Surg Res* 2011;97:705–11.
- [12] Ahlback S. Osteoarthritis of the knee. A radiographic investigation. *Acta Radiol Diagn* 1968;277(Suppl.):7–72.
- [13] Caton J, Deschamps G, Chambat P, Lerat JL, Dejour H. Patella infera. Apropos of 128 cases. *Rev Chir Orthop Rep Appar Mot* 1982;68:317–25.
- [14] Dejour H, Bonnin M. Tibial translation after anterior cruciate ligament rupture. Two radiological tests compared. *J Bone Joint Surg Br* 1994;76:745–9.
- [15] Mainard D. Les greffes osseuses. Cahiers d'enseignement de la SOFCOT. Paris: Elsevier; 2009. p. 297–322.
- [16] Amendola A, Bonasia DE. Results of high tibial osteotomy: review of the literature. *Int Orthop* 2010;34:155–60.
- [17] Bonasia DE, Governale G, Spolaore S, Rossi R, Amendola A. High tibial osteotomy. *Curr Rev Musculoskelet Med* 2014;7:292–301.
- [18] DeMeo PJ, Johnson EM, Chiang PP, Flamm AM, Miller MC. Midterm follow-up of opening-wedge high tibial osteotomy. *Am J Sports Med* 2010;38:2077–84.
- [19] Duivenvoorden T, Brouwer RW, Baan A, Bos PK, Reijman M, Bierma-Zeinstra SMA, et al. Comparison of closing-wedge and opening-wedge high tibial osteotomy for medial compartment osteoarthritis of the knee: a randomized controlled trial with a six-year follow-up. *J Bone Joint Surg Am* 2014;96:1425–32.
- [20] Harris JD, McNeilan R, Siston RA, Flanagan DC. Survival and clinical outcome of isolated high tibial osteotomy and combined biological knee reconstruction. *The Knee* 2013;20:154–61.
- [21] Hernigou P, Flouzat Lachaniette C, Delambre J, Guissou I, Dahmani O, Ibrahim Ouali M, et al. Full weight bearing and dynamisation with Limmed® locked plate fixation accelerates bone regeneration in the volume of opening wedge high tibial osteotomy. *Int Orthop* 2015;39:1295–300.
- [22] Laprade RF, Spiridonov SI, Nystrom LM, Jansson KS. Prospective outcomes of young and middle-aged adults with medial compartment osteoarthritis treated with a proximal tibial opening wedge osteotomy. *Arthrosc J Arthrosc Relat Surg Off Publ Arthrosc Assoc N Am Int Arthrosc Assoc* 2012;28:354–64.
- [23] Saragaglia D, Blaysat M, Inman D, Mercier N. Outcome of opening wedge high tibial osteotomy augmented with a Biosorb® wedge and fixed with a plate and screws in 124 patients with a mean of ten years follow-up. *Int Orthop* 2011;35:1151–6.
- [24] Smith TO, Sexton D, Mitchell P, Hing CB. Opening- or closing-wedged high tibial osteotomy: a meta-analysis of clinical and radiological outcomes. *The Knee* 2011;18:361–8.
- [25] Hernigou P, Roussignol X, Flouzat-Lachaniette CH, Filippini P, Guissou I, Poignard A. Opening wedge tibial osteotomy for large varus deformity with Ceraver resorbable beta tricalcium phosphate wedges. *Int Orthop* 2010;34:191–9.
- [26] Schallberger A, Jacobi M, Wahl P, Maestretti G, Jakob RP. High tibial valgus osteotomy in unicompartmental medial osteoarthritis of the knee: a retrospective follow-up study over 13–21 years. *Knee Surg Sports Traumatol Arthrosc* 2011;19:122–7.
- [27] Niinimäki TT, Eskelinen A, Mann BS, Junnila M, Ohtonen P, Leppilahti J. Survivorship of high tibial osteotomy in the treatment of osteoarthritis of the knee: Finnish registry-based study of 3195 knees. *J Bone Joint Surg Br* 2012;94:1517–21.
- [28] Bode G, von Heyden J, Pestka J, Schmal H, Salzmänn G, Südkamp N, et al. Prospective 5-year survival rate data following open-wedge valgus high tibial osteotomy. *Knee Surg Sports Traumatol Arthrosc Knee Surg Sports Traumatol Arthrosc* 2015;23:1949–55.
- [29] Martin R, Birmingham TB, Willits K, Litchfield R, Lebel M-E, Giffin JR. Adverse event rates and classifications in medial opening wedge high tibial osteotomy. *Am J Sports Med* 2014;42:1118–26.
- [30] Miller BS, Downie B, McDonough EB, Wojtyls EM. Complications after medial opening wedge high tibial osteotomy. *Arthroscopy* 2009;25:639–46.
- [31] Ramanoudjame M, Vandenbussche E, Baring T, Solignac N, Augereau B, Gregory T. Fibular nonunion after closed-wedge high tibial osteotomy. *Orthop Traumatol Surg Res* 2012;98:863–7.
- [32] Darnis A, Villa V, Debette C, Lustig S, Servien E, Neyret P. Vascular injuries during closing-wedge high tibial osteotomy: A cadaveric angiographic study. *Orthop Traumatol Surg Res* 2014;100:891–4.
- [33] Hernigou P, Goutallier D. [Outcome of the femoropatellar joint in osteoarthritic genu varum after tibial wedge osteotomy for angulation. 10 to 13 year regression]. *Rev Chir Orthop Réparatrice App Mot* 1987;73:43–8.
- [34] Lootvoet L, Massinon A, Rossillon R, Himmer O, Lambert K, Ghose JP. [Upper tibial osteotomy for gonarthrosis in genu varum. Apropos of a series of 193 cases reviewed 6 to 10 years later]. *Rev Chir Orthop Rép App Mot* 1993;79:375–84.
- [35] Bonasia DE, Dettoni F, Sisto G, Blonna D, Marmotti A, Bruzzone M, et al. Medial opening wedge high tibial osteotomy for medial compartment overload/arthritis in the varus knee: prognostic factors. *Am J Sports Med* 2014;42:690–8.
- [36] Amzallag J, Pujol N, Maqdes A, Beaufils P, Judet T, Catonne Y. Patellar height modification after high tibial osteotomy by either medial opening-wedge or lateral closing-wedge osteotomies. *Knee Surg Sports Traumatol Arthrosc* 2013;21:255–9.
- [37] Ducat A, Sariali E, Lebel B, Mertil P, Hernigou P, Flecher X, et al. Posterior tibial slope changes after opening- and closing-wedge high tibial osteotomy: a comparative prospective multicenter study. *Orthop Traumatol Surg Res* 2012;98:68–74.
- [38] Jacobi M, Villa V, Reischl N, Demey G, Goy D, Neyret P, et al. Factors influencing posterior tibial slope and tibial rotation in opening wedge high tibial osteotomy. *Knee Surg Sports Traumatol Arthrosc* 2015;23:2762–8.
- [39] Lee YS, Lee SB, Oh WS, Kwon YE, Lee BK. Changes in patellofemoral alignment do not cause clinical impact after open-wedge high tibial osteotomy. *Knee Surg Sports Traumatol Arthrosc* 2014. <http://dx.doi.org/10.1007/s00167-014-3349-y>.
- [40] Schröter S, Lobenhoffer P, Mueller J, Ihle C, Stöckle U, Albrecht D. Changes of patella position after closed and open wedge high tibial osteotomy: review of the literature. *Orthop* 2012;41(186):188–94.
- [41] El Amrani MH, Lévy B, Scharlycki S, Asselineau A. Patellar height relevance in opening-wedge high tibial osteotomy. *Orthop Traumatol Surg Res* 2010;96:37–43.
- [42] Lash NJ, Feller JA, Batty LM, Wasiak J, Richmond AK. Bone grafts and bone substitutes for opening-wedge osteotomies of the knee: a systematic review. *Arthroscopy* 2015;31:720–30.
- [43] Cho SW, Kim DH, Lee GC, Lee SH, Park SH. Comparison between autogenous bone graft and allogeneous cancellous bone graft in medial open wedge high tibial osteotomy with 2-year follow-up. *Knee Surg Relat Res* 2013;25:117–25.
- [44] Kuremsky MA, Schaller TM, Hall CC, Roehr BA, Masonis JL. Comparison of autograft vs allograft in opening-wedge high tibial osteotomy. *J Arthroplasty* 2010;25:951–7.
- [45] Hooper NM, Schouten R, Hooper GJ. The outcome of bone substitute wedges in medial opening high tibial osteotomy. *Open Orthop J* 2013;7:373–7.
- [46] Gouin F, Yaouanc F, Waast D, Melchior B, Delecir J, Passuti N. Open wedge high tibial osteotomies: Calcium-phosphate ceramic spacer versus autologous bonegraft. *Orthop Traumatol Surg Res* 2010;96:637–45.
- [47] Ganji R, Omidvar M, Izadfar A, Alavina SM. Opening wedge high tibial osteotomy using tibial wedge allograft: a case series study. *Eur J Orthop Surg Traumatol Orthopédie Traumatol* 2013;23:111–4.
- [48] Jung KA, Lee SC, Ahn NK, Hwang SH, Nam CH. Radiographic healing with hemispherical allogeneic femoral head bone grafting for opening-wedge high tibial osteotomy. *Arthroscopy* 2010;26:1617–24.
- [49] Lee SC, Jung KA, Nam CH, Jung SH, Hwang SH. The short-term follow-up results of open wedge high tibial osteotomy with using an Aescula open wedge plate and an allogeneic bone graft: the minimum 1-year follow-up results. *Clin Orthop Surg* 2010;2:47–54.
- [50] Santic V, Tudor A, Sestan B, Legovic D, Sirola L, Rakovac I. Bone allograft provides bone healing in the medial opening high tibial osteotomy. *Int Orthop* 2010;34:225–9.